

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec*

EPA Reg. No. 62719-375

For Use as a Protectant Fungicide for Control of Powdery Mildew Diseases on Cherry

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec* fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

Refer to product label for General Use Precautions and Restrictions and Mixing instructions.

| Disease | Recommended Spray Interval (days) | Rate of Quintec (fl oz/acre) [†] |
|--|-----------------------------------|---|
| powdery mildew (<i>Podosphaera clandestina</i>) | 10 - 14 | 7 |

[†] When applied as a dilute spray, maintain the per acre use rate.

Application Timing: Apply Quintec before visible symptoms of powdery mildew appear. Quintec has no curative properties and will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative spray or, if multiple diseases are present, following application of a broad-spectrum fungicide. Under low disease conditions (as per a predictive modeling system), minimum label use rates per application can be used. Maximum label rates are recommended for severe or threatening disease conditions.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- **Maximum Application Rate:** Do not apply more than 7 fl oz of Quintec per acre per application or more than 35 fl oz of Quintec per acre per calendar year.
- Do not make more than 5 applications of Quintec per calendar year.
- **Minimum Spray Interval:** 7 days.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

*Trademark of Dow AgroSciences LLC

R136-001

EPA-accepted 09/30/03

Initial printing