
PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

SUPPLEMENTAL LABEL FOR SWITCH 62.5WG

FUNGICIDE

EPA Reg. No. 100-953

For control of diseases on berries, pistachio and watercress

Active Ingredients:

Cyprodinil (CAS No. 121552-61-2)	37.5%
Fludioxonil (CAS No. 131341-86-1)	25.0%
<hr/>	
<i>Other Ingredients:</i>	37.5%
<i>Total:</i>	100.0%

Switch 62.5WG is a water-dispersible granule.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

SCP 953A-S1 0303

USE DIRECTIONS FOR SPECIFIC CROPS

Rotational Crop Restrictions: Do not plant any other crop for a period of 12 months unless Switch 62.5WG is registered for that use.

Crop	Disease	Rate oz./Acre	Application Timing	Notes
<p>Berry Crop Grouping:</p> <p>Bushberries (blueberries, currants, gooseberries, elderberry, huckleberry, as well as other cultivars and hybrids of these).</p> <p>Caneberries (blackberries, raspberries, and/or cultivars and/or hybrids of these).</p> <p>Juneberry Lingonberry Salal</p>	<p>Mummy Berry (<i>Monilinia vaccinii-corymbosi</i>)</p> <p>Anthracnose (<i>Colletotrichum spp.</i>)</p> <p>Alternaria Fruit Rot (<i>Alternaria tenuissima</i>)</p> <p>Phomopsis (<i>Phomopsis vaccinii</i>)</p> <p>Botrytis Fruit Rot (<i>Botryis cinerea</i>)</p>	<p>11-14</p>	<p>Make the first application during early bloom. A second application should be made 7 to 10 days later. Additional applications can be made at 7-10 day intervals if conditions remain favorable for disease development.</p> <p>Make no more than two (2) sequential applications before using another registered fungicides. Switch 62.5WG may be applied in an alternating or blocking program.</p>	<p>Pre-harvest interval is 0 days.</p> <p>Do not apply more than 56 ounces of product per acre per year.</p>

Crop	Disease	Rate oz./Acre	Application Timing	Notes
Pistachio	<p>Botrytis (<i>Botrytis</i>)</p> <p>Alternaria (<i>Alternaria</i>)</p>	11-14	<p>Make the first application during early bloom. A second application should be made 14 days later. Additional applications can be made at 14 day intervals if conditions remain favorable for disease development.</p> <p>Make no more than two (2) sequential applications before using another registered fungicide. Switch 62.5WG may be applied in an alternating or blocking program.</p>	<p>Do not apply Switch 62.5WG within 7 days of harvest. (7 day PHI)</p> <p>Do not apply more than 56 ounces of product per acre per year.</p>
Strawberries	Gray Mold (<i>Botrytis cinerea</i>)	11-14	<p>Begin application at or before bloom and continue on a 7 to 10-day interval.</p> <p>Make no more than two (2) sequential applications before using another registered fungicide. Switch 62.5WG may be applied in an alternating or blocking program .</p>	<p>Preharvest interval is 0-days.</p> <p>Do not exceed 56 ounces of product per year per acre.</p>
Watercress	<p>Cercospora leafspot (<i>Cercospora spp.</i>)</p> <p>Sclerotinia white mold (<i>Sclerotinia spp.</i>)</p> <p>Rhizoctonia rot (<i>Rhizoctonia solani</i>)</p>	11-14	<p>Begin applications when disease first appears and continue on a 7-10 day interval.</p> <p>Make no more than two (2) sequential applications before using another registered fungicides. Switch 62.5WG may be applied in an alternating or blocking program.</p>	<p>Pre-harvest interval is 0 days.</p> <p>Do not apply more than 56 ounces of product per acre per year.</p>

Switch™ and the Syngenta logo are trademarks of a Syngenta Group
Company
U.S. Patent No.5,153,200

© 2003 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP953A-S1 0303

SWI 62.5WG 953A-S1 0303 supp label - nsp – 3-17-03